

The Role of ISASS in Evolving the Spine Code Landscape: Lumbar Discogenic Pain Receives Specific ICD-10-CM Code

Morgan P. Lorio, Hansen A. Yuan, Douglas P. Beall, Jon E. Block and Gunnar B.J. Andersson

Int J Spine Surg published online 26 July 2024

<https://www.ijssurgery.com/content/early/2024/07/26/8622>

This information is current as of July 31, 2024.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

The Role of ISASS in Evolving the Spine Code Landscape: Lumbar Discogenic Pain Receives Specific ICD-10-CM Code

MORGAN P. LORIO, MD¹; HANSEN A. YUAN, MD²; DOUGLAS P. BEALL, MD³; JON E. BLOCK, PhD⁴; AND GUNNAR B.J. ANDERSSON, MD, PhD⁵

¹Advanced Orthopedics, Altamonte Springs, FL, USA; ²SUNY Upstate University, Syracuse, NY, USA; ³Comprehensive Specialty Care, Edmond, OK, USA; ⁴Consultant, San Francisco, CA, USA; ⁵Consultant, Scottsdale, AZ, USA

Keywords: discogenic, pain, lumbar, spine, degeneration

On behalf of the International Society for the Advancement of Spine Surgery (ISASS), we are pleased to announce that the National Center for Health Statistics has granted and issued specific International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) codes for lumbosacral discogenic pain associated with degenerative disc disease (Table).¹ This action represents a significant advancement in our ability to differentiate, classify, and precisely diagnose chronic back pain syndromes. Lumbar and lumbosacral discogenic pain will now be appropriately included within the ICD-10-CM rubric along with other sources of back pain (eg, neurocompressive, facetogenic, sacroiliac, vertebrogenic, etc). The new codes will take effect on 1 October 2024.

ISASS has been at the forefront in identifying the importance of discogenic pain and advocating for specific diagnostic coding.² Indeed, the intervertebral disc has been the most intensively studied and characterized structure in the human spine. While countless research groups have contributed to our rich knowledge about degeneration of the intervertebral disc, several key individuals and an event were instrumental in our current perception of lumbar discogenic pain. First, known as the original “needle jockeys,” the esteemed triumvirate of Professors Charles Aprill, Nikolai Bogduk, and Richard Derby contributed an enormous body of seminal experimental research isolating physiological, biochemical, and morphological characteristics of the disc as potent pain generators.³ Second, the development and clinical evaluation of intradiscal electrothermal therapy 2 decades ago provided renewed impetus to examine the anatomical features of internal disc disruption of

the annulus fibrosus.⁴ While the technology proved to be only modestly efficacious and did not enjoy commercial success, it spurred interest in the development of novel treatments aimed at disc degeneration.⁵

Under the new ICD-10-CM codes, symptomatic lumbar discogenic disease is diagnosed clinically by axial midline back pain, pain with flexion, sitting intolerance, positive provocative test with sustained hip flexion, absence of motor/sensor/reflex change, and elective discography.^{6,7} Discogenic back pain can occur with or without nonradicular/nonsciatic referred leg pain in a sclerotomal distribution.² We believe these codes will provide several advantages:

- **Precision in diagnosis:** Health care providers will have clearer criteria for diagnosing discogenic low back pain (LBP), allowing for more precise identification of the underlying pathology. This can guide treatment decisions and improve patient outcomes by ensuring that interventions are targeted to the specific cause of the pain.
- **Tailored treatment approaches:** With a better understanding of the specific nature of discogenic LBP, clinicians can tailor treatment approaches to individual patients. This may involve a combination of conservative therapies, minimally invasive procedures, and surgical interventions, depending on the severity and characteristics of the condition.
- **Innovation and research:** The availability of standardized codes can stimulate innovation and research in the field of spinal interventions. Researchers may be motivated to develop new techniques, devices, and therapies aimed at addressing discogenic LBP more effectively

Table. Current and new International Classification of Diseases, 10th Revision, Clinical Modification diagnostic codes for discogenic pain.

Current Code: Description	New Code: Description
Lumbar Region	
· M51: Thoracic, thoracolumbar, and lumbosacral intervertebral disc disorders	· M51.360: Other intervertebral disc degeneration, lumbar region with lumbar discogenic back pain only
· M51.3: Other thoracic, thoracolumbar and lumbosacral intervertebral disc degeneration	· M51.361: Other intervertebral disc degeneration, lumbar region with lower extremity pain only
· M51.36: Other intervertebral disc degeneration, lumbar region	· M51.362: Other intervertebral disc degeneration, lumbar region with discogenic back pain and lower extremity pain
	· M51.369: Other intervertebral disc degeneration, lumbar region without mention of lumbar back pain or lower extremity pain
Lumbosacral Region	
· M51.37: Other intervertebral disc degeneration, lumbosacral region	· M51.370: Other intervertebral disc degeneration, lumbosacral region with discogenic back pain only
	· M51.371: Other intervertebral disc degeneration, lumbosacral region with lower extremity pain only
	· M51.372: Other intervertebral disc degeneration, lumbosacral region with discogenic back pain and lower extremity pain
	· M51.379: Other intervertebral disc degeneration, lumbosacral region without mention of lumbar back pain or lower extremity pain

if they know there is a clearer pathway for reimbursement and utilization.

- **Expanded access to advanced therapies:** Reimbursement pathways for disc restoration, regeneration, and arthroplasty can increase patient access to advanced therapies for discogenic LBP, such as regenerative medicine techniques, motion-preserving implants, and other cutting-edge interventions previously underutilized due to reimbursement barriers.
- **Improved documentation and communication:** Standardized coding for discogenic LBP enhances documentation and communication between health care providers, insurers, and regulatory agencies. This can streamline administrative processes, facilitate accurate billing and reimbursement, and ensure that patients receive appropriate care based on established guidelines and protocols.

The ICD remains the standard (and most important) classification and diagnostic tool for epidemiology, health management, and clinical purposes. It is used globally as the foundation for the identification of health trends and statistics. Appropriately and accurately classifying patients with lumbar discogenic pain represents a significant, albeit long overdue, step forward in spinal care.

REFERENCES

1. 2025 ICD-10-CM. CMS.gov. 2025. <https://www.cms.gov/medicare/coding-billing/icd-10-codes/2025-icd-10-cm>.
2. Lorio MP, Beall DP, Calodney AK, Lewandrowski KU, Block JE, Mekhail N. Defining the patient with lumbar discogenic pain:

real-world implications for diagnosis and effective clinical management. *J Pers Med*. 2023;13(5):821. doi:10.3390/jpm13050821

3. Bogduk N, Aprill C, Derby R. Lumbar discogenic pain: state-of-the-art review. *Pain Med*. 2013;14(6):813–836. doi:10.1111/pme.12082

4. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine*. 1995;20(17):1878–1883. doi:10.1097/00007632-199509000-00007

5. Lorio MP, Tate JL, Myers TJ, Block JE, Beall DP. Perspectives on Intradiscal therapies for lumbar discogenic pain: state of the science, knowledge gaps, and Imperatives for clinical adoption. *J Pain Res*. 2024;17:1171–1182. doi:10.2147/JPR.S441180

6. Bogduk N. On the definitions and physiology of back pain, referred pain, and radicular pain. *Pain*. 2009;147(1–3):17–19. doi:10.1016/j.pain.2009.08.020

7. Kallewaard JW, Terheggen MAMB, Groen GJ, et al. 15. discogenic low back pain. *Pain Pract*. 2010;10(6):560–579. doi:10.1111/j.1533-2500.2010.00408.x

Funding: No funding was received for this article.

Disclosures: The authors report no conflicts of interest or disclosures.

Editor's Note: Dr Lorio is the current US president of ISASS. Dr Yuan is editor in chief emeritus of the International Journal of Spine Surgery and was a founding member of ISASS. Dr Andersson is a past president of ISASS.

Corresponding Author: Jon E. Block, Consultant, 2210 Jackson St, Suite 401, San Francisco, CA 94115, USA; jb@drjonblock.com

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2024 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.